

Eckmann-Hilton and the Hopf Fibration

Raymond Baker

University of Colorado Boulder

morphismz.github.io

HoTT/UF 2024

April 3rd, 2024

The Goal

And some reasons to care

The Goal: Construct the Hopf fibration $\text{hpf} : \mathbb{S}^3 \rightarrow \mathbb{S}^2$ using the Eckmann-Hilton argument.

And some reasons to care:

- 1 Simple description of the generator of $\pi_3(\mathbb{S}^2)$. From the fiber sequence of hpf .
- 2 Ditto the generator of $\pi_4(\mathbb{S}^3)$. From the Freudenthal suspension theorem.
- 3 $\pi_4(\mathbb{S}^3)$ has order *at most* 2. From Syllepsis.

The Plan

- 1 Use Eckmann-Hilton to construct $eh : \Omega^3(\mathbb{S}^2)$. This is equivalent to a map $hpf : \mathbb{S}^3 \rightarrow \mathbb{S}^2$.
- 2 Characterize the fiber as \mathbb{S}^1 by generalizing ideas from Kraus and Von Raumer's "Path Spaces of Higher Inductive Types".

The Eckmann-Hilton Argument

Eckmann-Hilton

For $\alpha, \beta : \Omega^2(X)$, we have $\text{EH}(\alpha, \beta) : \alpha \cdot \beta = \beta \cdot \alpha$

But where does this identification come from?

Where does Path Concatination come from?

Fix a pointed type (X, \bullet) and consider $\text{Id}_\bullet : X \rightarrow U$.

A loop $p : \Omega(X)$ induces:

$$\text{tr}^{\text{Id}_\bullet}(p) : \Omega(X) \simeq \Omega(X)$$

This is path concatenation:

for $q : \Omega(X)$ we have:

$$\text{tr}(p)(q) = q \cdot p.$$

Where does Eckmann-Hilton come from?

Up one dimension:

a 2-loop $\alpha : \Omega^2(X, \bullet)$ induces:

$$\mathrm{tr}^2(\alpha) : \mathrm{id}_{\Omega(X)} \sim \mathrm{id}_{\Omega(X)}$$

This is Eckmann-Hilton:

for $\beta : \Omega^2(X)$, we have:


$$\mathrm{nat}[\mathrm{tr}^2(\alpha)](\beta) = \mathrm{EH}(\alpha, \beta)$$

(modulo coherence paths)

A formula for $\text{tr}^2(\alpha)$

Computing $\text{tr}^2(\alpha) : \text{id}_{\Omega(X)} \sim \text{id}_{\Omega(X)}$

$$\text{tr}^2(\alpha) = \text{whisker}_\alpha = \lambda(p).\text{refl}_p \star \alpha$$



$$\text{tr}^2(\alpha)(\text{refl}_\bullet) = \alpha$$

The naturality condition of $\text{tr}^2(\alpha) : \text{id}_{\Omega(X)} \sim \text{id}_{\Omega(X)}$

For $\beta : \Omega^2(X)$:

$$\begin{array}{ccc} \text{refl.} & \xrightarrow{\text{id}(\beta)} & \text{refl.} \\ \text{tr}^2(\alpha)(\text{refl}) \downarrow & \text{nat-}[\text{tr}^2(\alpha)](\beta) & \downarrow \text{tr}^2(\alpha)(\text{refl}) \\ \text{refl.} & \xrightarrow{\text{id}(\beta)} & \text{refl.} \end{array}$$

Plus coherence paths, this defines

$$\text{EH}(\alpha, \beta) : \alpha \cdot \beta = \beta \cdot \alpha$$

.

Eckmann-Hilton in \mathbb{S}^2

$$\text{EH}(\text{surf}_2, \text{surf}_2) : \text{surf}_2 \cdot \text{surf}_2 = \text{surf}_2 \cdot \text{surf}_2$$

The type of this identification is equivalent to $\Omega^3(\mathbb{S}^2)$.

The Eckmann-Hilton 3-loop

Define $\text{eh} : \Omega^3(\mathbb{S}^2)$ as the image of $\text{EH}(\text{surf}_2, \text{surf}_2)$ under said equivalence.

See [agda-unimath](#) for more.

The map hpf

The 3-loop eh is equivalent to a map, the Hopf fibration:

$$\text{hpf} : \mathbb{S}^3 \rightarrow \mathbb{S}^2$$

Define a map $\text{hpf} : \mathbb{S}^3 \rightarrow \mathbb{S}^2$ by \mathbb{S}^3 -induction:

$$\text{hpf}(\text{base}_3) := \text{base}_2$$

$$\text{hpf}(\text{surf}_3) := \text{eh}$$

The Universal Property of the Family of Fibers

Fix a pointed map $h : A \rightarrow B$. Then:

Heuristic

$\text{fib}_h(b_0)$ is like the loop space of B with extra identifications freely generated by the map h .

The Universal Property of the Family of Fibers

We have an induced type family $\text{fib}_h \circ h : A \rightarrow U$.

This family always comes equipped with a section:

$$\lambda(a).(a, \text{refl}_{h(a)}) : (a : A) \rightarrow \text{fib}_h \circ h(a)$$

called a lift of h to fib_h .

The Wild Category of Families with Lifts

And the Universal Property of the Family of Fibers

Wild Category of Families with Lifts

Objects: families $P : B \rightarrow U$ equipped with a lift $(a : A) \rightarrow P \circ h(a)$

Maps: families of maps $(b : B) \rightarrow P(b) \rightarrow Q(b)$ that preserve the lift

Universal Property of fib_h

The family fib_h with its canonical lift is initial in this wild category.

Proof: follows from the standard equivalence $A \simeq \sum_{b:B} \text{fib}_h(b)$.
Formalized in [agda-unimath](#)

Loop Spaces are a Special Case

If $A \equiv \text{unit}$ and $h : \text{unit} \rightarrow B$ defined by $h(\star) \equiv b_0$:

$$((a : \text{unit}) \rightarrow P \circ h(a)) \simeq P(b_0)$$

So fib_h is the initial type family equipped with a point over b_0

Specializing the Universal Property

Let $A \equiv \mathbb{S}^3$, $B \equiv \mathbb{S}^2$ and $h \equiv \text{hpf}$.

Then fib_{hpf} is the initial:

family over \mathbb{S}^2

point $u : \text{fib}_{\text{hpf}}(\text{base}_2)$

identification $t : \text{tr}^3(\text{eh})(u) = \text{refl}_u^2$

The latter identification is equivalent to an identification

$$\text{tr}^3(\text{EH}(\text{surf}_2, \text{surf}_2))(u) = \text{refl}_{\text{tr}^2(\text{surf}_2 \cdot \text{surf}_2)}(u)$$

Specializing the Universal Property

fib_{hpf} is the initial:

family over \mathbb{S}^2

point $u : \text{fib}_{\text{hpf}}(\text{base}_2)$

identification $t : \text{tr}^3(\text{EH}(\text{surf}_2, \text{surf}_2))(u) = \text{refl}_{\text{tr}^2(\text{surf}_2 \cdot \text{surf}_2)}(u)$

Interlude, descent data of \mathbb{S}^2

A type family P over \mathbb{S}^2 is equivalent to:

Descent data of \mathbb{S}^2

a type X , the value of $P(\text{base}_2)$

a 2-automorphism $\text{id}_X \sim \text{id}_X$, the transport $\text{tr}^2(\text{surf}_2)$

A Characterization of fib_{hpf}

Then fib_{hpf} is the initial data:

type F

2-automorphism $H : \text{id}_F \sim \text{id}_F$

point $u : F$

identification $\text{tr}^3(\text{EH}(\text{surf}_2, \text{surf}_2))(u) = \text{refl}_{\text{tr}^2(\text{surf}_2 \cdot \text{surf}_2)}(u)$

Eckmann-Hilton in the Universe

For $P : X \rightarrow U$ with $u : P(\bullet)$ and $\alpha, \beta : \Omega^2(X, \bullet)$:

$$\begin{array}{ccc}
 \text{tr}^2(\alpha \cdot \beta)(u) & \xrightarrow{\text{tr}^2\text{-concat}_{\alpha, \beta}} & \text{tr}^2(\alpha)(u) \cdot \text{tr}^2(\beta)(u) \\
 \text{tr}^3(\text{EH}(\alpha, \beta))(u) \Bigg| & \text{tr}^3\text{-EH} & \Bigg| \text{nat}[\text{tr}^2(\alpha)](\text{tr}^2(\beta)(u)) \\
 \text{tr}^2(\beta \cdot \alpha)(u) & \xrightarrow{\text{tr}^2\text{-concat}_{\beta, \alpha}} & \text{tr}^2(\beta)(u) \cdot \text{tr}^2(\alpha)(u)
 \end{array}$$

Proof: See [agda-unimath](#)

A Characterization of fib_{hpf}

So fib_{hpf} is the initial data:

type F

2-automorphism $H : \text{id}_F \sim \text{id}_F$

point $u : F$

identification $\text{nat-}[\text{tr}^2(\text{surf}_2)](\text{tr}^2(\text{surf}_2)(u)) = \text{refl}_{\text{tr}^2(\text{surf}_2)(u)} \cdot \text{tr}^2 \text{surf}_2(u)$

A Characterization of fib_{hpf}

Finally, fib_{hpf} is the initial data:

type F

point $u : F$

2-automorphism $H : \text{id}_F \sim \text{id}_F$

identification $\text{nat-}H(H(u)) = \text{refl}_{H(u)} \cdot H(u)$

The Fiber is \mathbb{S}^1

Want $F \simeq \mathbb{S}^1$

Two approaches:

- 1 Using a HIT and directly constructing an equivalence
- 2 Show \mathbb{S}^1 is initial in the wild category of F -algebras

Using a HIT

In cubical agda: thanks to Tom Jack

In Book HoTT: possible ...

In agda-unimath (and other common HoTT repos): not possible

F -algebras

Give a definition of the wild category of F -algebras

Then show $\mathrm{hom}_{F\text{-alg}}(\mathbb{S}^1, X)$ is contractible for every F -algebra X .

\mathbb{S}^1 forms an F -algebra

type - \mathbb{S}^1

2-automorphism - L

point - b_1

identification - $\text{defn}_L : \text{nat-L}(L(b_1)) = \text{refl}_{\text{loop}} \cdot \text{loop}$

Morphisms of F -algebras

Consider an F -algebra (X, K, x_0, p)

A morphism of F -algebras $(\mathbb{S}^1, L, b_1, \text{defn}_L) \rightarrow (X, K, x_0, p)$ comprises:

- 1 $g : \mathbb{S}^1 \rightarrow X$
- 2 $G : g \cdot_l L \sim K \cdot_l g$
- 3 $g_0 : g(b_1) = x_0$
- 4 t , a witness that “ defn_L is sent to q ”

$\text{hom}_{F\text{-alg}}(\mathbb{S}^1, X) \simeq \text{unit}$

a map: $(g : \mathbb{S}^1 \rightarrow X, G : g \cdot_l L \sim K \cdot_r g, g_0 : g(b_1) = x_0, t)$

g is equivalent to $g(b_1) : X$ and $g(\text{loop}) : \Omega(X, x)$.

$(g(b_1), g_0)$ is a contractible pair.

G is equivalent to $G(b) : g(\text{loop}) = K(g(b_1))$ and $\text{nat-}G(\text{loop})$.

$(g(\text{loop}), G)$ is a contractible pair.

Claim: $\text{nat-}G(\text{loop})$ and t form a contractible pair.

Fiber Sequence and the Calculation $\pi_3(\mathbb{S}^2)$

We now have a fiber sequence $\mathbb{S}^1 \rightarrow \mathbb{S}^3 \xrightarrow{\text{hpf}} \mathbb{S}^2$

Consequences:

It follows that $\Omega^3(\text{hpf}) : \Omega^3(\mathbb{S}^3) \simeq \Omega^3(\mathbb{S}^2)$

So $\text{eh} : \Omega^3(\mathbb{S}^2)$ generates $\pi_3(\mathbb{S}^2) \cong \mathbb{Z}$

$\pi_4(\mathbb{S}^3)$ has order ≤ 2

The Generator of $\pi_4(\mathbb{S}^3)$

eh_{surf_3} generates $\pi_4(\mathbb{S}^3)$

Proof: Freudenthal + functions preserve eh .

$\pi_4(\mathbb{S}^3)$ has order ≤ 2

The square of eh_{surf_3} is trivial.

Proof: Syllepsis (see Sojakova)

Future Work

- ① non-triviality of $\pi_4(\mathbb{S}^3)$ (a full calculation of $\pi_4(\mathbb{S}^3)$)
- ② Adapting the James construction and Wärrn's Zig Zag Construction
- ③ Higher Hopf Fibrations and Higher Coherences

Non-Triviality of $\pi_4(\mathbb{S}^3)$

Suffices to find a family $B : \Omega(\mathbb{S}^3) \rightarrow U$ such that

$$\text{nat}[\text{tr}^2(\text{surf}_3)](\text{tr}^2(\text{surf}_3)(u))$$

is non-trivial, for some $u : B(\text{refl})$

It would follow $\pi_4(\mathbb{S}^3) \cong \mathbb{Z}/2\mathbb{Z}$.

Adapting James and Zig Zag

the worst part of the proof: the recursive HIT, showing its \mathbb{S}^1

This is a familar problem to those characterizing loop spaces.

The solution (for certain cases):

suspension: the James construction

pushouts: Zig Zag construction

A hope: versions of these constructions for general fibers (already in the literature?)

Higher Hopf Fibrations and their Coherences

The higher Hopf fibrations $\mathbb{S}^7 \rightarrow \mathbb{S}^4$ and $\mathbb{S}^{15} \rightarrow \mathbb{S}^8$ should also arise from higher coherences.

The E_4 coherence, corresponding to $\mathbb{S}^7 \rightarrow \mathbb{S}^4$, was constructed by Sojakova.

E_n and Descent over \mathbb{S}^n

$\text{surf}_n : \Omega^n(\mathbb{S}^n)$ induces an n -automorphism of $\Omega(\mathbb{S}^n)$

the E_n coherence is the $(n - 1)$ -dimensional naturality condition this.

easy to calculate for $n = 1, 2$. I've calculated this for $n = 3$ with much trouble. The case for $n \geq 4$ needs a motivated approach

Acknowledgements

- My Advisor: Professor Jonathan Wise, CU Boulder
- Tom Jack, for many helpful discussions
- Egbert Rijke, Fredrik Bakke, Vojtěch Štěpančík, and the agda-unimath community

The End

Questions? Comments?